

RESEARCH

Open Access



# Adherence to 24-h movement behavior guidelines and psychosocial functioning in young children: a longitudinal analysis

Rachael W. Taylor<sup>1\*</sup>, Jillian J. Haszard<sup>2</sup>, Dione Healey<sup>3</sup>, Kim A. Meredith-Jones<sup>1</sup>, Barry J. Taylor<sup>4</sup> and Barbara C. Galland<sup>4</sup>

## Abstract

**Background:** A recent paradigm shift has highlighted the importance of considering how sleep, physical activity and sedentary behaviour work together to influence health, rather than examining each behaviour individually. We aimed to determine how adherence to 24-h movement behavior guidelines from infancy to the preschool years influences mental health and self-regulation at 5 years of age.

**Methods:** Twenty-four hour movement behaviors were measured by 7-day actigraphy (physical activity, sleep) or questionnaires (screen time) in 528 children at 1, 2, 3.5, and 5 years of age and compared to mental health (anxiety, depression), adaptive skills (resilience), self-regulation (attentional problems, hyperactivity, emotional self-control, executive functioning), and inhibitory control (Statue, Head-Toes-Knees-Shoulders task) outcomes at 5 years of age. Adjusted standardised mean differences (95% CI) were determined between those who did and did not achieve guidelines at each age.

**Results:** Children who met physical activity guidelines at 1 year of age (38.7%) had lower depression (mean difference [MD]: -0.28; 95% CI: -0.51, -0.06) and anxiety (MD: -0.23; 95% CI: -0.47, 0.00) scores than those who did not. At the same age, sleeping for 11–14 h or having consistent wake and sleep times was associated with lower anxiety (MD: -0.34; 95% CI: -0.66, -0.02) and higher resilience (MD: 0.35; 95% CI: 0.03, 0.68) scores respectively. No significant relationships were observed at any other age or for any measure of self-regulation. Children who consistently met screen time guidelines had lower anxiety (MD: -0.43; 95% CI: -0.68, -0.18) and depression (MD: -0.36; 95% CI: -0.62, -0.09) scores at 5. However, few significant relationships were observed for adherence to all three guidelines; anxiety scores were lower (MD: -0.42; 95% CI: -0.72, -0.12) in the 20.2% who adhered at 1 year of age, and depression scores were lower (MD: -0.25; 95% CI: -0.48, -0.02) in the 36.7% who adhered at 5 years of age compared with children who did not meet all three guidelines.

**Conclusions:** Although adherence to some individual movement guidelines at certain ages throughout early childhood was associated with improved mental health and wellbeing at 5 years of age, particularly reduced anxiety and depression scores, there was little consistency in these relationships. Future work should consider a compositional approach to 24-h time use and how it may influence mental wellbeing.

**Trial registration:** ClinicalTrials.gov number [NCT00892983](https://clinicaltrials.gov/ct2/show/study/NCT00892983)

\*Correspondence: [rachael.taylor@otago.ac.nz](mailto:rachael.taylor@otago.ac.nz)

<sup>1</sup> Department of Medicine, University of Otago, PO Box 56, Dunedin, New Zealand

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Keywords:** Child, Physical activity, Sleep, Screen time, 24-h movement behaviors, Mental health, Wellbeing, Psychosocial functioning

## Background

The World Health Organization defines mental health as a state of wellbeing and effective functioning in which an individual recognises his or her own abilities, is resilient to the stresses of life, and is able to make a positive contribution to his or her community [1]. Mental health difficulties can be expressed at a very early age, grouped into two broad categories of internalizing (e.g. anxiety, depression) and externalizing (e.g. hyperactivity, aggression) symptomatology. A hallmark feature of these difficulties is the ability to self-regulate, that is, exercise control over one's thoughts, feelings, and behaviors. Self-regulation can be demonstrated in several ways, including an ability to focus and regulate attention or suppress a thought or action (inhibitory control), and in childhood, is associated with concurrent and later levels of achievement, interpersonal skills, mental health, and healthy living [2].

While many adverse environmental and social factors contribute to poor mental health in children, positive lifestyle factors such as getting enough good quality sleep, undertaking regular physical activity, and limiting time spent sedentary are also considered to be important [3–5]. However, relatively little work has examined how these behaviors influence mental health in preschool-aged children, a time of rapid growth and development. Recent reviews have highlighted the scarcity of literature (particularly that which is longitudinal in nature) and the inconsistent findings observed in the predominantly cross-sectional literature linking mental health and wellbeing outcomes with sleep [6], physical activity [7], or sedentary time [8] when each behaviour is examined in isolation.

However, a paradigm shift has occurred in the last few years which emphasises that we also need to be investigating these behaviors simultaneously because while each behavior can independently affect health, they also interact in ways that may not be apparent if studied individually [9, 10]. Consequently, many countries have introduced 24-h movement behavior guidelines for young children in recognition of this issue [11–14]. However, few studies have examined adherence to 24-h guidelines in relation to mental health in preschool-aged children [15–19], a time when sleep, physical activity and sedentary behaviors patterns are changing rapidly. The three cross-sectional studies reported some significant relationships between adherence to guidelines and improvements in behaviors,

emotions [16], social cognition [15], or psychosocial or physical health [19]. However, in all studies, effect sizes were small and other measures evaluated showed no relationship with health outcomes [15, 16] or no effect on total health [19]. The two longitudinal studies both demonstrated no evidence of association between compliance to individual or combined guidelines and a variety of psychosocial outcomes [17, 18].

However, there are some methodological issues with these studies. While some of these studies have used objective measures of physical activity [16–18], only one assessed the impact of additional guidelines relating to activity intensity [16]. Only one of these studies also measured sleep objectively [16], although they did not examine the effect of additional guidelines relating regularly in sleep patterns. Although two studies assessed relationships between movement behaviors and mental health longitudinally [17, 18], repeated measures of movement behaviors over this time of rapid developmental change [20] have not been undertaken.

Therefore, the aim of this study was to determine how adherence to 24-h movement behavior guidelines from infancy (<2 years) to the preschool years (2–5 years) is related to mental health and self-regulation at 5 years of age. It is hypothesized that greater time spent asleep or active, and less time spent sedentary will lead to improved mental health and a greater ability to self-regulate behaviour when children reach the age of 5 years.

## Methods

This analysis uses data from children participating in the Prevention of Overweight in Infancy (POI) randomised controlled trial undertaken in Dunedin, New Zealand. POI investigated whether anticipatory guidance for parents about infant sleep, dietary intake, and physical activity reduced excessive weight gain, compared to usual care. Anticipatory guidance is that given by the health care provider to assist parents to understand how their children grow and develop. The intervention commenced in late pregnancy and concluded when the child was two years of age [21], with follow-up measurements obtained at 3.5 and 5 years of age (just prior to starting school – children in New Zealand typically start school on or just after their fifth birthday) [22]. As the intervention did not produce differences in physical activity, sedentary behavior, or sleep [23–25], these data have been analysed using the entire cohort with appropriate adjustment for randomisation group.

As detailed information on the original trial is available in both the study protocols [21, 22] and published findings [24, 25], only brief details are provided here. The original intervention was approved by the Lower South Ethics Committee (LRS/12/08/063) and the follow-up study by the University of Otago Human Ethics Committee (12/274) and was powered to detect differences in body mass index (BMI) between intervention groups. Written informed consent was obtained from the parent/guardian of all child participants. All mothers who had booked into the single maternity hospital (>97% of all births) serving the population of Dunedin, New Zealand were invited to participate when in mid to late pregnancy (May 2009 to December 2010). A 58% response rate yielded a final sample size of 802 primiparous (47.6%) and multiparous (52.4%) mothers. Women were randomized to one of four study groups by the study biostatistician, within 6 strata depending on area level deprivation (3 levels) and parity (2 levels), using a block size of 12. Allocation was concealed using opaque, pre-sealed envelopes. All anthropometric assessments and accelerometry analyses were performed by researchers blinded to group allocation. Demographic information obtained at baseline (late pregnancy) included maternal age, education, ethnicity, self-reported pre-pregnancy height and weight, and area level deprivation calculated using NZDep 2013 [26] (a deprivation score calculated from census data which reflects the extent of material and social deprivation, and ranges from 1 (least deprived) to 10 (most deprived)). Information on infant gestational age, sex and birth weight was obtained from hospital records. Anthropometric measurements were obtained by trained measurers following standard protocols at all subsequent time points [27]. Duplicate measures of weight (Tanita WB-100 MA/WB-110 MA) and height (Harpenden stadiometer, Holtain Ltd, UK) were obtained with children wearing underwear/light clothing [22]. Body mass index (BMI) z-scores were determined using the World Health Organization growth standards [28], with overweight defined as a BMI z-score  $\geq 85^{\text{th}}$  but  $< 95^{\text{th}}$  percentile, and obesity as BMI  $\geq 95^{\text{th}}$  percentile.

*Sleep and physical activity* were assessed with children wearing Actical (Mini-Mitter, Bend, OR) accelerometers 24-h a day for up to 7 days at 1, 2, 3.5 and 5 years of age. The accelerometers were worn around the waist on elastic belts and initialized using 15 s epochs. Data were cleaned and scored using an automated script developed in MATLAB (MathWorks, Natick, MA, USA) [29]. This count-scaled algorithm estimates sleep onset (start of the first 15 continuous minutes of sleep preceded by 5 min of awake) and offset (last of 15 continuous minutes of sleep followed by 5 min of awake) specific to each individual for each day. Night sleep duration is calculated as the

duration of time between sleep onset and offset, excluding time awake [30]. Day-time sleep was determined at 1 and 2 years of age by the automated Matlab script which defines naps during daytime wake periods (9am to 5 pm) as at least 30 min of continuous minutes of sleep, preceded by 5 min of awake [29]. All sleep data reported here refer to total sleep duration (night-time sleep and naps, where relevant, combined). Non-wear time, sedentary time, and time in light to vigorous physical activity is then determined during awake hours only. Non-wear time was defined as at least 20 min of consecutive zeros [31]. We defined sedentary time as 0–6 counts (per 15 s), and light-to-vigorous physical activity (LMVPA) as  $\geq 7$  counts [32, 33]. A day was considered valid if the participant had at least eight hours of wear time outside of sleep time. Participants were only included in these analyses at each age if they had at least three valid days of data. Data from all valid days were averaged, weighted depending on whether the day was a weekend day or a weekday.

*Screen time* was assessed at 1, 2, and 5 years of age via interviewer-administered questionnaires with the parent/guardian (>99% mothers). At 1 and 2 years of age, parents indicated the frequency per week, and the number of minutes each time, their child ‘usually’ watched television, videos or DVDs. We expanded the number of media types measured when the children were 5 years of age to include television (free to air, cable, and online streaming), DVDs or videos, computers for games (not school-work), passive games consoles (including handheld), and mobile phones or tablets for playing games. Parents indicated the frequency per week and duration each time their child spent watching/using each category of media use resulting in a total duration per day (minutes). Screen time information was not collected at 3.5 years of age.

*Restraint* was measured at 1 and 2 years of age only, where parents indicated the number of times over the past week, and the length of time each session, the infant was placed in a car seat or stroller.

#### **Adherence to guidelines**

Adherence to the Canadian guidelines for the early years (0–4 years) [11] was calculated using all available data at each age, including the 5 year old measurements given most (79.9%) children were measured before their 5th birthday (mean age 4.98 (0.4) years). Level 1 adherence was defined as: *physical activity*; at least 180 min of light-to-vigorous physical activity (LMVPA) each day (average over all relevant days of measurement) at 1, 2, 3.5 and 5 years, *sedentary*; no screen time (1- and 2-year-old children) or  $\leq 60$  min per day (5-year-old children), and *sleep*; receiving 11–14 h (1- and 2-year-old children) or 10–13 h (3.5 and 5-year-old children) every 24 h. We also calculated Level 2 adherence guidelines as follows:

*physical activity*; including at least 60 min of energetic play (defined as moderate-to-vigorous activity, MVPA) in the 3.5 and 5-year-old physical activity recommendations, *sedentary*; restricting the use of restraint (such as in car seats or strollers) for no more than an hour at a time at 1 and 2 years only, and *sleep*; having consistent bed and wake times (calculated as a within-person SD for sleep onset and offset of < 30 min) at all four ages.

### Mental health outcomes at 5 years of age

Mental wellbeing and resilience were measured by parental report using the Anxiety (13 items, covering levels of everyday worry, fear, and perfectionism  $\alpha=0.82$ ), Depression (11 items, covering mood and negative cognitions,  $\alpha=0.77$ ), and Resilience (12 items, covering adjustment to setbacks or unexpected events and social functioning,  $\alpha=0.82$ ) subscales of the Behavioral Assessment System for Children (BASC-2) 2–5 year old scale [34]. The BASC-2 is a well-validated and normed scale designed to assess wide-ranging areas of child functioning as rated by parents. Self-regulation was measured using the BASC-2 Hyperactivity (11 items, covering activity level, impulsivity, and self-control,  $\alpha=0.79$ ), Attentional Problems (6 items, covering the ability to attend and follow instructions,  $\alpha=0.80$ ), Emotional Self-Control (6–8 items, covering the ability manage negative emotions and frustrations,  $\alpha=0.79$ ), and Executive Functioning (10–13 items, covering inhibitory control, distractibility, and emotional reactivity,  $\alpha=0.79$ ) subscales; as well as objective measures of Inhibitory control using the ‘Statue’ component of the NeuroPSYchological Assessment (NEPSY-2) [35] which measures difficulty with inhibitory control, and the Head-Toes-Knees-Shoulders task [36]. The NEPSY-2 is a test battery assessing numerous areas of neuropsychological functioning. It is well-normed, reliable, and appropriate for use with 5-year-old children. The Head-Toes-Knees-Shoulders task [36] is a measure of behavioral regulation and inhibition, which determines the ability of a child to follow opposing instructions (e.g. touch the head when directed to touch the toes) that is designed for 3–7 year old children.

### Statistical analyses

All statistical analysis was undertaken in Stata 16.1 (StataCorp, Texas). Only participants who had completed the 5-year assessments of mental health, had at least one assessment of movement behaviors, and had complete data on covariates (randomised group, sex, maternal education, household deprivation, primiparity, and BMI z-score) were included in these analyses. Demographics were described for the subsamples used at each age. As all mental health assessments were scored

using different scales, all scores were standardised to be in units of standard deviations (SD), with the exception of the heads-toes-knees-shoulder task, which had right-skewed data that were not normalized by log-transformation and is described using the median, and the 25<sup>th</sup> and 75<sup>th</sup> percentiles. For ease of interpretation, three scales were reverse scored compared to the BASC-2 scoring protocol so that high scores represented more of the behavior (Resilience, Emotional self-control, and Executive functioning).

Mean differences with 95% confidence intervals (CI) in the standardised scores between those who met the guideline and those who did not meet the guideline were estimated using linear regression models, adjusted for the covariates listed previously. Quantile regression for the median was used for the heads-toes-knees-shoulder assessment. Estimates were not calculated if the group sizes were too small – this was decided to be the case if one of the groups was less than 10% of the sample. Associations are judged on the size of the mean difference as well as the 95% CI, and not solely on whether the p-value passes the threshold of ‘statistical significance’ (<0.05) [37]. Here we use the convention that an effect size greater than 0.2SD has the potential to be meaningful [38]. Residuals of models were plotted and visually assessed for heteroskedasticity and normality.

### Results

Although 528 participants had at least one measurement of movement behaviors at any age and mental health outcomes at 5 years of age, the number of participants included at each age ranged from 256 (2 years of age) to 382 (5 years of age). The demographics at baseline across these samples were broadly comparable, with 44–51% of all infants being female and first-born. A high proportion of mothers had tertiary education, and a lower proportion of children came from homes with higher levels of deprivation than is nationally expected (Table 1). Participants included in these analyses ( $n=528$ ) did not differ in terms of baseline demographics from those not followed up ( $n=274$ ) for infant sex, maternal ethnicity, parity, or household deprivation [39]. Table 2 presents the possible minimum and maximum scores for each index, along with the mean (SD) scores for each of the mental health measures at 5 years of age.

Table 3 presents the standardised mean difference (95% CI) in mental wellbeing scores according to whether children met each of the physical activity, sedentary time, and sleep guidelines, and combinations thereof, at each age. The 38.7% of children who participated in at least 180 min of light-to-vigorous physical activity (LMVPA) each day at 1 year of age had lower scores for depression (mean difference [MD]: -0.28;

**Table 1** Demographics of the analysed sample at 1, 2, 3.5, and 5 years of age<sup>a</sup> (*n* = 528)

|  | Age (years) |             |             |             |             |            |
|--|-------------|-------------|-------------|-------------|-------------|------------|
|  | All         | 1           | 2           | 3.5         | 5           |            |
| N  | 528         | 315         | 256         | 305         | 382         |            |
| Maternal age at child's birth, mean (SD) years | 32.6 (4.6)  | 32.9 (4.5)  | 32.9 (4.4)  | 33.0 (4.5)  | 32.8 (4.5)  |            |
| Primiparous, n (%)                             | 249 (47.2)  | 159 (50.5)  | 112 (43.8)  | 134 (43.9)  | 170 (44.5)  |            |
| Maternal ethnicity, n (%)                      | European    | 462 (87.5)  | 282 (89.5)  | 229 (89.5)  | 270 (88.5)  | 337 (88.2) |
|  | Māori       | 21 (4.0)    | 11 (3.5)    | 6 (2.3)     | 11 (3.6)    | 14 (3.7)   |
|  | Other       | 45 (8.5)    | 22 (7.0)    | 21 (8.2)    | 24 (7.9)    | 31 (8.1)   |
| Maternal tertiary education, n (%)             | 359 (68.0)  | 224 (71.1)  | 180 (70.3)  | 205 (67.2)  | 258 (67.5)  |            |
| Household deprivation                          | Low         | 196 (37.1)  | 116 (36.8)  | 104 (40.6)  | 122 (40.0)  | 142 (37.2) |
|  | Medium      | 231 (43.8)  | 142 (45.1)  | 106 (41.4)  | 130 (42.6)  | 168 (44.0) |
|  | High        | 101 (19.1)  | 57 (18.1)   | 46 (18.0)   | 53 (17.4)   | 72 (18.9)  |
| Infant sex, n (%) female                       | 256 (48.5)  | 155 (49.2)  | 117 (45.7)  | 146 (47.9)  | 191 (50.0)  |            |
| BMI z-score at 5 years of age, mean (SD)       | 0.46 (0.89) | 0.41 (0.84) | 0.43 (0.87) | 0.47 (0.87) | 0.48 (0.89) |            |

<sup>a</sup> Analysed sample determined by those with data on mental wellbeing at 5 years of age, all demographics reported, and physical activity data at other ages

**Table 2** Mental wellbeing, self-regulation and inhibitory control measures at 5 years of age (*n* = 528)

| Outcome   | Subscale                  | n   | Possible minimum and maximum scores | Mean (SD)               |
|---|---------------------------|-----|-------------------------------------|-------------------------|
| Mental wellbeing and adaptive skills <sup>a</sup> | Depression                | 528 | 0 to 33                             | 7.2 (3.5)               |
|   | Anxiety                   | 528 | 0 to 39                             | 8.8 (4.6)               |
|   | Resilience                | 528 | 0 to 36                             | 25.6 (4.8)              |
| Self-regulation <sup>a</sup>                      | Attentional problems      | 528 | 0 to 18                             | 6.4 (2.5)               |
|   | Hyperactivity             | 528 | 0 to 33                             | 9.5 (4.0)               |
|   | Emotional self-control    | 528 | 0 to 24                             | 18.6 (3.1)              |
|   | Executive functioning     | 528 | 0 to 39                             | 27.0 (4.2)              |
| Inhibitory control <sup>b</sup>                   | Statue                    | 463 | 0 to 30                             | 17.1 (8.8)              |
|   | Head-toes-knees-shoulders | 463 | 0 to 60                             | 27 (7, 40) <sup>c</sup> |

<sup>a</sup> Mental wellbeing, adaptive skills, and self-regulation all assessed using the Behavioral Assessment System for children (BASC-2) [34], with higher scores reflecting higher levels of construct

<sup>b</sup> Inhibitory control assessed using laboratory based tasks. The 'Statue' assessment from the NeuroPSYchological Assessment (NEPSY-2) [35] was scored from 0 (movements throughout assessment) to 30 (no movements). The Heads-toes-knees-shoulder [36] assessment was scored from 0 (all incorrect with no self-correction) to 60 (all correct)

<sup>c</sup> median (25th, 75th percentile) reported for Head-toes-knees-shoulders assessment because of right-skew

95% CI: -0.51, -0.06) and anxiety (MD: -0.23; 95%CI: -0.47, 0.00) and higher scores for resilience (MD: 0.23; 95%CI: -0.01, 0.46) at 5 years of age than children who did not meet this guideline. Corresponding effect sizes could not be calculated at 2, 3.5 or 5 years of age because of ceiling effects in compliance (94.1 to 98.4% of children). However, fewer children met PA guidelines at these ages once the requirement to have 60 min of energetic play was included within the 180 min of LMVPA (54.1 to 81.2%), but few comparisons at this age met our requirement for a difference of at least 0.20 to be potentially meaningful, with no differences being statistically significant.

Greater consistency was observed in relation to compliance with limiting screen time, in that scores for anxiety and depression were often about one-third of a standard deviation lower in children who met screen time behavior guidelines than those who did not. Effects on anxiety (MD: -0.43; 95% CI: -0.68 to -0.18) and depression (MD: -0.36; 95% CI: -0.62, -0.09) were particularly marked for those who met the screen time guidelines on at least two occasions. By contrast, restricting screen time at any single age appeared to have little effect on resilience scores, unless children had consistently limited screen use (MD: 0.24; 95% CI: -0.02, 0.50). Almost all children were not restrained in carseats or prams etc. for more than one

**Table 3** Meeting 24-h movement behavior guidelines through early childhood and mental wellbeing (depression and anxiety) and adaptive skills (resilience) at five years ( $n = 528$ )

|   | n   | N (%)<br>who met<br>guideline | Depression at 5 years of age<br>Standardised mean difference (95% CI) <sup>a</sup> for those who met the guideline compared<br>to those who did not | Anxiety at 5 years of age<br>Standardised mean difference (95% CI) <sup>a</sup> for those who met the guideline compared<br>to those who did not | Resilience at 5 years of age<br>Standardised mean difference (95% CI) <sup>a</sup> for those who met the guideline compared<br>to those who did not |
|---|-----|-------------------------------|---|--|---|
| Physical activity: $\geq 180$ min/day LMVPA <sup>b</sup>  |     |                               |   |  |   |
| At 1 year of age  | 315 | 122 (38.7)                    | <b>-0.28 (-0.51, -0.06)</b>   | <b>-0.23 (-0.47, 0.00)</b>   | 0.23 (-0.01, 0.46)  |
| Physical activity: $\geq 180$ min/day LMVPA including 60 min energetic play <sup>c</sup>  |     |                               |   |  |   |
| At 3.5 years of age   | 305 | 187 (61.3)                    | -0.04 (-0.28, 0.20)   | -0.04 (-0.28, 0.20)  | 0.05 (-0.17, 0.28)  |
| At 5 years of age   | 382 | 310 (81.2)                    | -0.11 (-0.38, 0.16)   | -0.20 (-0.46, 0.06)  | 0.21 (-0.04, 0.47)  |
| At both 3.5 and 5 years of age  | 266 | 144 (54.1)                    | -0.12 (-0.37, 0.13)   | -0.15 (-0.40, 0.10)  | 0.17 (-0.07, 0.40)  |
| Sedentary time: no screen time or no more than 60 min/day <sup>d</sup>  |     |                               |   |  |   |
| At 1 year of age  | 305 | 191 (62.6)                    | -0.15 (-0.38, 0.08)   | <b>-0.30 (-0.54, -0.06)</b>  | 0.12 (-0.11, 0.36)  |
| At 2 years of age   | 220 | 35 (15.9)                     | -0.29 (-0.66, 0.07)   | -0.17 (-0.55, 0.21)  | 0.13 (-0.25, 0.52)  |
| At 5 years of age   | 377 | 151 (40.1)                    | <b>-0.34 (-0.56, -0.13)</b>   | -0.13 (-0.34, 0.09)  | 0.18 (-0.03, 0.39)  |
| At 2 or more time points  | 311 | 83 (26.7)                     | <b>-0.43 (-0.68, -0.18)</b>   | <b>-0.36 (-0.62, -0.09)</b>  | 0.24 (-0.02, 0.50)  |
| Sedentary time: not being restrained for > 60 min at a time <sup>e</sup>  |     |                               |   |  |   |
| At 1 year of age  | 305 | 273 (89.5)                    | -0.14 (-0.51, 0.22)   | 0.00 (-0.38, 0.39)   | 0.07 (-0.31, 0.44)  |
| Sedentary time: no screen time or no > 60 min AND not being restrained for > 60 min at a time   |     |                               |   |  |   |
| At 1 year of age  | 305 | 176 (57.7)                    | -0.12 (-0.34, 0.11)   | <b>-0.29 (-0.53, -0.06)</b>  | 0.11 (-0.12, 0.34)  |
| Sleep: 11 – 14 h or 10 – 13 h including naps <sup>f</sup>   |     |                               |   |  |   |
| At 1 year of age  | 307 | 258 (84.0)                    | -0.07 (-0.37, 0.23)   | <b>-0.34 (-0.66, -0.02)</b>  | 0.13 (-0.18, 0.43)  |
| At 5 years of age   | 370 | 325 (87.8)                    | 0.18 (-0.14, 0.50)  | 0.00 (-0.31, 0.32)   | -0.12 (-0.43, 0.19)   |
| Sleep: consistent sleep and wake times <sup>g</sup>   |     |                               |   |  |   |
| At 1 year of age  | 307 | 56 (18.2)                     | -0.20 (-0.49, 0.07)   | -0.13 (-0.43, 0.17)  | 0.25 (-0.04, 0.54)  |
| At 2 years of age   | 254 | 67 (26.4)                     | 0.11 (-0.17, 0.40)  | 0.13 (-0.16, 0.43)   | -0.06 (-0.36, 0.23)   |
| At 3.5 years of age   | 398 | 59 (19.8)                     | 0.13 (-0.16, 0.43)  | 0.15 (-0.14, 0.44)   | -0.04 (-0.32, 0.24)   |
| At 5 years of age   | 370 | 84 (22.7)                     | -0.21 (-0.46, 0.04)   | -0.08 (-0.32, 0.17)  | 0.08 (-0.16, 0.33)  |
| At 2 or more time points  | 380 | 58 (15.3)                     | -0.14 (-0.43, 0.14)   | -0.11 (-0.40, 0.18)  | 0.04 (-0.24, 0.32)  |
| Sleep: both sleep guidelines (duration and consistency)   |     |                               |   |  |   |
| At 1 year of age  | 307 | 42 (13.7)                     | -0.25 (-0.57, 0.07)   | -0.19 (-0.53, 0.16)  | <b>0.35 (0.03, 0.68)</b>  |
| At 2 years of age   | 254 | 64 (25.2)                     | 0.11 (-0.18, 0.40)  | 0.10 (-0.19, 0.40)   | -0.06 (-0.36, 0.24)   |
| At 3.5 years of age   | 298 | 56 (18.8)                     | 0.22 (-0.08, 0.51)  | 0.21 (-0.09, 0.51)   | -0.11 (-0.40, 0.18)   |
| At 5 years of age   | 370 | 77 (20.8)                     | -0.20 (-0.46, 0.05)   | -0.03 (-0.29, 0.22)  | 0.03 (-0.22, 0.29)  |
| At 2 or more time points  | 380 | 48 (12.6)                     | -0.13 (-0.44, 0.17)   | 0.01 (-0.30, 0.33)   | -0.04 (-0.34, 0.26)   |
| All three guidelines: $\geq 180$ min/day LMVPA, no screen time or no more than 60 min, and 11 – 14 h or 10 – 13 h including naps <sup>h</sup> |     |                               |   |  |   |
| At 1 year of age  | 297 | 60 (20.2)                     | -0.19 (-0.47, 0.09)   | <b>-0.42 (-0.72, -0.12)</b>  | 0.28 (-0.01, 0.57)  |
| At 2 years of age   | 218 | 33 (15.1)                     | -0.23 (-0.60, 0.14)   | -0.13 (-0.52, 0.26)  | 0.09 (-0.31, 0.49)  |
| At 5 years of age   | 365 | 134 (36.7)                    | <b>-0.25 (-0.48, -0.02)</b>   | -0.11 (-0.33, 0.11)  | 0.12 (-0.11, 0.34)  |

<sup>a</sup> Assessed using the Behavioral Assessment System for children (BASC-2) [34]

<sup>a</sup> Mean difference (95% CI) estimated using regression models adjusted for randomised group, sex, maternal education, household deprivation, primiparous, and BMI z-score at 5 years of age, in units of standard deviations. Greater scores indicate higher levels of depression, anxiety, or resilience

<sup>b</sup> Proportions who met the first physical activity guideline at 2, 3.5, and 5 years of age were too high to allow estimation of mean differences (94.1, 97.1, and 98.4%, respectively)

<sup>c</sup> Second physical activity guideline applies only to children aged 3 – 5 years

<sup>d</sup> No screen time (1 – 2 years) or 60 min maximum screen time (3 – 5 years). Not measured at 3.5 years

<sup>e</sup> Not being restrained in a car seat or stroller for > 60 min at a time, on at least 3 days a week (1 – 2 years). The proportion who met this guideline at 2 years of age was too high to allow estimation of mean differences (96.8%). Not measured at 3.5 years

<sup>f</sup> 11 – 14 h (1 – 2 years) or 10 – 13 h (3 – 5 years). The proportion who met this guideline at 2 and 3.5 years of age and at 2 or more time points was too high to allow estimation of mean differences (92.9%, 95.6%, and 92.6% respectively)

<sup>g</sup> Having a consistent sleep time or consistent wake time defined as within-person standard deviation of < 30 min

<sup>h</sup> The proportion who met this guideline at two or more time points was too low to allow estimation of mean differences (9.7%)

Bold figures denote  $P < 0.05$

hour at a time, with no apparent relationship with scores for mental wellbeing (Table 3).

Table 3 also indicates that while approximately one-quarter of the effect sizes for sleep compliance in relation to mental wellbeing met our threshold of potentially meaningful, only two were statistically significant. Children who slept for 11–14 h at 1 year of age had lower scores for anxiety at 5 years of age (MD: -0.34; 95% CI: -0.66, -0.02) and those who also had consistent wake and sleep times at this age had higher resilience scores at 5 years (MD: 0.35; 95% CI: 0.03, 0.68). Otherwise, meeting sleep guidelines, whether examined in terms of duration (level 1) or consistency in bed and wake times (level 2), or both, had relatively little impact on wellbeing outcomes at 5 years of age.

Children who met guidelines for all three behaviors at 1, 2, or 5 years of age had lower scores for depression when they were aged 5 years, although the difference was only statistically significant in the cross-sectional analysis at 5 years of age (MD: -0.25; 95% CI: -0.48, -0.02). Effect sizes for anxiety and resilience were more variable. The 20.2% of children who adhered to guidelines for all three behaviors at 1 year of age had lower scores for anxiety at age 5 (MD: -0.42; 95% CI: -0.72, -0.12), but meaningful effect sizes were not observed at other ages. No statistically significant differences were observed for resilience, although compliance at the age of 1 was associated with potentially better resilience at age 5 (MD: 0.28; 95% CI: -0.01, 0.57, Table 3).

Adherence to physical activity or sleep guidelines at any age had little relationship to measures of self-regulation at 5 years of age (Table 4). While a few individual results met our criteria for effect sizes that were potentially meaningful, these were not consistent over time and none were statistically significant. Meeting screen time guidelines was not associated with measures of self-regulation at any individual age. However, children who met screen time guidelines on at least two points (26.7%) had significantly lower scores for hyperactivity (MD: -0.26; 95% CI: -0.52, -0.01), and significantly higher scores for emotional self-control (MD: 0.35; 95% CI: 0.11, 0.59) and executive functioning (MD: 0.28; 95% CI: 0.03, 0.53) (Table 4). Not being restrained in carseats and strollers for more than 1 h at a time at the age of 1 year was associated with lower scores for attentional problems (MD: -0.25; 95% CI: -0.60, 0.09) and hyperactivity scores (MD: -0.37; 95% CI: -0.74, -0.00), but not emotional self-control or executive functioning at 5 years of age. Similar relationships could not be examined at two years of age because almost all (96.8%) of children met this guideline (Table 4).

Relatively few significant differences were observed in levels of inhibitory control according to adherence

to guidelines in preschoolers (Table 5). Higher levels of inhibitory control, as indicated by higher Statue scores, were observed for those who met physical activity guidelines at 3.5 (MD: 0.25; 95% CI: -0.01, 0.52) or both 3.5 and 5 years of age (MD: 0.24; 95% CI: 0.00, 0.49). Neither compliance with screen time guidelines or those targeting limits to time spent restrained appeared related to measures of inhibitory control. Similarly, although children who achieved sufficient sleep duration at the age of 5 showed higher levels of inhibitory control (MD: 0.36; 95% CI: 0.05, 0.68), as did those who had consistent sleep/wake times at 1 year of age (MD: 0.37; 95% CI: 0.06, 0.69), no other sleep comparisons were significant. Only 1 of 27 comparisons were significant for the Heads-toes-knees-shoulder task as a measure of behavioral regulation and inhibition, with scores being higher for children who were physically active at 3.5 years of age (median difference: 9.0; 95% CI: 1.4, 16.5). Overall, meeting all three behavior guidelines did not appear related to either of these measures of inhibitory control at any age.

## Discussion

Our longitudinal study demonstrates that achieving appropriate levels of sleep and physical activity while limiting sedentary time appears to have relatively little impact on measures of mental health and wellbeing at five years of age. Overall, meeting individual physical activity or sleep guidelines did not appear consistently related to levels of anxiety or depression in preschool-aged children, nor did this predict improved self-regulation or inhibitory control at this age. By contrast, scores for anxiety and depression were often about one-third of a standard deviation lower in children who met sedentary behavior guidelines compared with those who did not at any age. Children who met guidelines for all three behaviors tended to have lower scores for anxiety at 5 years of age, whereas findings for depression and resilience were more variable. Meeting all guidelines was not related to significant differences in self-regulation or inhibitory control at any age examined.

It is difficult to compare our findings with the literature given so few studies have examined this question in early life. Comparisons are also limited by the wide range in outcomes that have been investigated, with little consistency across the relevant studies. A recent review in older children and adolescents has suggested favourable associations between meeting all three recommendations and improved mental health when compared with meeting none of the recommendations [40]. While this appears somewhat in contrast to the current study, these authors highlight that the evidence in general is weak, and in relation to each specific mental health indicator, was limited [40]. Data in younger children are scarce. A

**Table 4** Meeting 24-h movement behavior guidelines through early childhood and self-regulation at five years ( $n = 528$ )

|   | n   | N (%)<br>who met<br>guideline | Attentional<br>problems at 5 years<br>of age | Hyperactivity at<br>5 years of age | Emotional self-<br>control at 5 years<br>of age | Executive<br>functioning at<br>5 years of age |
|---|-----|-------------------------------|--|------------------------------------|---|---|
| Standardised mean difference (95% CI) <sup>a</sup> for those who met the guideline compared to those who did not                              |     |                               |  |                                    |   |   |
| Physical activity: $\geq 180$ min/day LMVPA <sup>b</sup>  |     |                               |  |                                    |   |   |
| At 1 year of age  | 315 | 122 (38.7)                    | -0.18 (-0.40, 0.04)                          | -0.11 (-0.35, 0.12)                | 0.18 (-0.03, 0.40)                              | 0.10 (-0.12, 0.33)                            |
| Physical activity: $\geq 180$ min/day LMVPA including 60 min energetic play <sup>c</sup>  |     |                               |  |                                    |   |   |
| At 3.5 years of age   | 305 | 187 (61.3)                    | 0.00 (-0.21, 0.20)                           | 0.16 (-0.07, 0.39)                 | 0.02 (-0.21, 0.24)                              | -0.17 (-0.39, 0.06)                           |
| At 5 years of age   | 382 | 310 (81.2)                    | -0.09 (-0.34, 0.15)                          | 0.15 (-0.11, 0.41)                 | 0.05 (-0.20, 0.31)                              | 0.02 (-0.23, 0.27)                            |
| At both 3.5 and 5 years of age  | 266 | 144 (54.1)                    | 0.01 (-0.21, 0.22)                           | 0.22 (-0.01, 0.46)                 | 0.04 (-0.19, 0.26)                              | -0.11 (-0.34, 0.12)                           |
| Sedentary time: no screen time or no > 60 minutes <sup>d</sup>  |     |                               |  |                                    |   |   |
| At 1 year of age  | 305 | 191 (62.6)                    | -0.09 (-0.31, 0.13)                          | -0.15 (-0.38, 0.09)                | 0.12 (-0.09, 0.34)                              | 0.07 (-0.15, 0.30)                            |
| At 2 years of age   | 220 | 35 (15.9)                     | -0.04 (-0.39, 0.31)                          | 0.00 (-0.37, 0.37)                 | 0.12 (-0.22, 0.47)                              | 0.13 (-0.25, 0.50)                            |
| At 5 years of age   | 377 | 151 (40.1)                    | -0.14 (-0.34, 0.06)                          | -0.07 (-0.28, 0.15)                | 0.17 (-0.03, 0.38)                              | 0.14 (-0.07, 0.35)                            |
| At 2 or more time points  | 311 | 83 (26.7)                     | -0.11 (-0.35, 0.14)                          | <b>-0.26 (-0.52, -0.01)</b>        | <b>0.35 (0.11, 0.59)</b>                        | <b>0.28 (0.03, 0.53)</b>                      |
| Sedentary time: not being restrained for > 60 min at a time <sup>e</sup>  |     |                               |  |                                    |   |   |
| At 1 year of age  | 305 | 273 (89.5)                    | -0.25 (-0.60, 0.09)                          | <b>-0.37 (-0.74, -0.00)</b>        | 0.14 (-0.21, 0.48)                              | 0.19 (-0.17, 0.55)                            |
| Sedentary time: no screen time or no > 60 min AND not being restrained for > 60 min at a time   |     |                               |  |                                    |   |   |
| At 1 year of age  | 305 | 176 (57.7)                    | -0.12 (-0.34, 0.09)                          | -0.15 (-0.38, 0.08)                | 0.10 (-0.11, 0.31)                              | 0.05 (-0.17, 0.28)                            |
| Sleep: 11 – 14 h or 10 – 13 h including naps <sup>f</sup>   |     |                               |  |                                    |   |   |
| At 1 year of age  | 307 | 258 (84.0)                    | -0.01 (-0.30, 0.28)                          | -0.05 (-0.36, 0.26)                | 0.07 (-0.21, 0.35)                              | 0.09 (-0.21, 0.39)                            |
| At 5 years of age   | 370 | 325 (87.8)                    | 0.14 (-0.15, 0.44)                           | 0.04 (-0.28, 0.35)                 | -0.03 (-0.34, 0.28)                             | -0.12 (-0.42, 0.18)                           |
| Sleep: consistent sleep and wake times <sup>g</sup>   |     |                               |  |                                    |   |   |
| At 1 year of age  | 307 | 56 (18.2)                     | -0.19 (-0.47, 0.08)                          | -0.04 (-0.33, 0.25)                | 0.21 (-0.06, 0.48)                              | 0.17 (-0.11, 0.45)                            |
| At 2 years of age   | 254 | 67 (26.4)                     | 0.09 (-0.18, 0.36)                           | 0.01 (-0.28, 0.30)                 | -0.12 (-0.40, 0.15)                             | -0.09 (-0.38, 0.20)                           |
| At 3.5 years of age   | 398 | 59 (19.8)                     | -0.04 (-0.30, 0.21)                          | 0.07 (-0.21, 0.34)                 | -0.14 (-0.41, 0.14)                             | 0.02 (-0.26, 0.29)                            |
| At 5 years of age   | 370 | 84 (22.7)                     | -0.18 (-0.41, 0.05)                          | -0.19 (-0.43, 0.06)                | 0.16 (-0.08, 0.40)                              | 0.13 (-0.10, 0.37)                            |
| At 2 or more time points  | 380 | 58 (15.3)                     | -0.19 (-0.46, 0.07)                          | -0.10 (-0.38, 0.18)                | 0.03 (-0.24, 0.30)                              | 0.15 (-0.13, 0.43)                            |
| Sleep: both sleep guidelines (duration and consistency)   |     |                               |  |                                    |   |   |
| At 1 year of age  | 307 | 42 (13.7)                     | -0.17 (-0.48, 0.14)                          | -0.07 (-0.40, 0.25)                | 0.22 (-0.08, 0.52)                              | 0.23 (-0.09, 0.55)                            |
| At 2 years of age   | 254 | 64 (25.2)                     | 0.05 (-0.23, 0.32)                           | -0.01 (-0.31, 0.28)                | -0.08 (-0.36, 0.20)                             | -0.06 (-0.36, 0.23)                           |
| At 3.5 years of age   | 298 | 56 (18.8)                     | 0.03 (-0.23, 0.29)                           | 0.13 (-0.15, 0.41)                 | -0.23 (-0.51, 0.06)                             | -0.06 (-0.34, 0.22)                           |
| At 5 years of age   | 370 | 77 (20.8)                     | -0.12 (-0.36, 0.12)                          | -0.17 (-0.42, 0.08)                | 0.17 (-0.08, 0.42)                              | 0.11 (-0.14, 0.35)                            |
| At 2 or more time points  | 380 | 48 (12.6)                     | -0.15 (-0.44, 0.14)                          | -0.07 (-0.37, 0.24)                | 0.00 (-0.30, 0.29)                              | 0.15 (-0.15, 0.45)                            |
| All three guidelines: $\geq 180$ min/day LMVPA, no screen time or no more than 60 min, and 11 – 14 h or 10 – 13 h including naps <sup>h</sup> |     |                               |  |                                    |   |   |
| At 1 year of age  | 297 | 60 (20.2)                     | -0.07 (-0.34, 0.20)                          | -0.08 (-0.37, 0.21)                | 0.17 (-0.09, 0.44)                              | 0.08 (-0.20, 0.37)                            |
| At 2 years of age   | 218 | 33 (15.1)                     | 0.05 (-0.31, 0.41)                           | 0.08 (-0.30, 0.46)                 | 0.02 (-0.34, 0.37)                              | 0.01 (-0.37, 0.40)                            |
| At 5 years of age   | 365 | 134 (36.7)                    | -0.05 (-0.26, 0.16)                          | -0.02 (-0.24, 0.20)                | 0.12 (-0.10, 0.34)                              | 0.07 (-0.15, 0.28)                            |

<sup>a</sup> Mean difference (95% CI) estimated using regression models adjusted for randomised group, sex, maternal education, household deprivation, primiparous, and BMI z-score at 5 years of age, in units of standard deviations

<sup>b</sup> Proportions who met the first physical activity guideline at 2, 3.5, and 5 years of age were too high to allow estimation of mean differences (94.1, 97.1, and 98.4%, respectively)

<sup>c</sup> Second physical activity guideline applies only to children aged 3 – 5 years

<sup>d</sup> No screen time (1 – 2 years) or 60 min maximum screen time (3 – 5 years). Not measured at 3.5 years

<sup>e</sup> Not being restrained in a car seat or stroller for > 60 min at a time, on at least 3 days a week (1 – 2 years). The proportion who met this guideline at 2 years of age was too high to allow estimation of mean differences (96.8%). Not measured at 3.5 years

<sup>f</sup> 11 – 14 h (1 – 2 years) or 10 – 13 h (3 – 5 years). The proportion who met this guideline at 2 and 3.5 years of age and at 2 or more time points was too high to allow estimation of mean differences (92.9, 95.6, and 92.6% respectively)

<sup>g</sup> Having a consistent sleep time or consistent wake time defined as within-person standard deviation of < 30 min

<sup>h</sup> The proportion who met this guideline at two or more time points was too low to allow estimation of mean differences (9.7%)

Bold figures denote  $P < 0.05$



**Table 5** Meeting 24-h movement behavior guidelines through early childhood and inhibitory control at five years ( $n = 463$ )

|   | n   | N (%)<br>who met<br>guideline | Statue at 5 years of age<br>Standardised mean difference (95% CI) <sup>a</sup> for<br>those who met the guideline compared to<br>those who did not | Heads-toes-knees-shoulder at 5 years of age<br>Median difference (95% CI) <sup>b</sup> for those who met<br>the guideline compared to those who did not |
|---|-----|-------------------------------|--|---|
| Physical activity: $\geq 180$ min/day LMVPA <sup>c</sup>  |     |                               |  |   |
| At 1 year of age  | 287 | 111 (38.7)                    | 0.03 (-0.22, 0.28)   | -2.3 (-10.0, 5.5)   |
| Physical activity: $\geq 180$ min/day LMVPA including 60 min energetic play <sup>d</sup>  |     |                               |  |   |
| At 3.5 years of age   | 293 | 180 (61.4)                    | 0.12 (-0.12, 0.35)   | <b>9.0 (1.4, 16.5)</b>  |
| At 5 years of age   | 372 | 304 (81.7)                    | 0.25 (-0.01, 0.52)   | 1.4 (-7.6, 10.4)  |
| At both 3.5 and 5 years of age  | 261 | 142 (54.4)                    | <b>0.24 (0.00, 0.49)</b>   | 7.3 (-0.9, 15.6)  |
| Sedentary time: no screen time or no > 60 min <sup>e</sup>  |     |                               |  |   |
| At 1 year of age  | 277 | 174 (62.8)                    | 0.08 (-0.18, 0.33)   | 1.3 (-6.0, 8.6)   |
| At 2 years of age   | 200 | 33 (16.5)                     | -0.21 (-0.59, 0.18)  | 4.0 (-7.6, 15.7)  |
| At 5 years of age   | 367 | 146 (39.8)                    | 0.15 (-0.07, 0.36)   | -1.0 (-7.5, 5.6)  |
| At 2 or more time points  | 296 | 80 (27.0)                     | 0.02 (-0.24, 0.28)   | 5.3 (-2.4, 13.1)  |
| Sedentary time: not being restrained for > 60 min at a time <sup>f</sup>  |     |                               |  |   |
| At 1 year of age  | 277 | 246 (88.8)                    | 0.06 (-0.33, 0.46)   | 5.7 (-5.5, 16.9)  |
| Sedentary time: no screen time or no > 60 min AND not being restrained for > 60 min at a time   |     |                               |  |   |
| At 1 year of age  | 277 | 159 (57.4)                    | 0.08 (-0.17, 0.33)   | 2.5 (-5.0, 10.0)  |
| Sleep: 11 – 14 h or 10 – 13 h including naps <sup>g</sup>   |     |                               |  |   |
| At 1 year of age  | 280 | 235 (83.9)                    | -0.06 (-0.40, 0.27)  | -5.9 (-15.7, 3.9)   |
| At 5 years of age   | 360 | 316 (87.8)                    | <b>0.36 (0.05, 0.68)</b>   | 9.5 (-1.1, 20.2)  |
| Sleep: consistent sleep and wake times <sup>h</sup>   |     |                               |  |   |
| At 1 year of age  | 280 | 51 (18.2)                     | <b>0.37 (0.06, 0.69)</b>   | 7.3 (-1.9, 16.6)  |
| At 2 years of age   | 233 | 60 (25.7)                     | 0.03 (-0.28, 0.33)   | -1.4 (-10.3, 7.5)   |
| At 3.5 years of age   | 286 | 58 (20.3)                     | 0.16 (-0.13, 0.45)   | 6.6 (-3.0, 16.2)  |
| At 5 years of age   | 360 | 82 (22.8)                     | 0.18 (-0.06, 0.43)   | 5.4 (-3.0, 13.9)  |
| At 2 or more time points  | 361 | 57 (15.8)                     | 0.12 (-0.17, 0.41)   | 4.5 (-4.5, 13.6)  |
| Sleep: both sleep guidelines (duration and consistency)   |     |                               |  |   |
| At 1 year of age  | 280 | 38 (13.6)                     | 0.30 (-0.05, 0.66)   | 4.6 (-5.5, 14.8)  |
| At 2 years of age   | 233 | 57 (24.5)                     | 0.02 (-0.29, 0.33)   | -1.0 (-9.9, 7.9)  |
| At 3.5 years of age   | 286 | 55 (19.2)                     | 0.19 (-0.10, 0.49)   | 5.6 (-4.1, 15.3)  |
| At 5 years of age   | 360 | 75 (20.8)                     | 0.19 (-0.07, 0.45)   | 2.9 (-5.8, 11.6)  |
| At 2 or more time points  | 361 | 47 (13.0)                     | 0.08 (-0.23, 0.40)   | -0.2 (-9.5, 9.2)  |
| All three guidelines: $\geq 180$ min/day LMVPA, no screen time or no > 60 min, and 11 – 14 h or 10 – 13 h including naps <sup>i</sup> |     |                               |  |   |
| At 1 year of age  | 270 | 53 (19.6)                     | -0.10 (-0.42, 0.22)  | 1.4 (-8.0, 10.7)  |
| At 2 years of age   | 198 | 31 (15.7)                     | -0.19 (-0.59, 0.21)  | 5.8 (-6.1, 17.7)  |
| At 5 years of age   | 355 | 129 (36.3)                    | 0.14 (-0.09, 0.36)   | 1.9 (-5.5, 9.2)   |

<sup>a</sup> Mean difference (95% CI) estimated using linear regression models adjusted for randomised group, sex, maternal education, household deprivation, primiparous, and BMI z-score at 5 years of age, in units of standard deviations

<sup>b</sup> Median difference (95% CI) estimated using quantile regression models adjusted for randomised group sex, maternal education, household deprivation, primiparous, and BMI z-score at 5 years of age, scores range from 0 (less inhibitory control) to 60 (more inhibitory control)

<sup>c</sup> Proportions who met the first physical activity guideline at 2, 3.5, and 5 years of age were too high to allow estimation of mean differences (93.6, 96.9, and 98.4%, respectively)

<sup>d</sup> Second physical activity guideline applies only to children aged 3 – 5 years

<sup>e</sup> No screen time (1 – 2 years) or 60 min maximum screen time (3 – 5 years). Not measured at 3.5 years

<sup>f</sup> Not being restrained in a car seat or stroller for > 60 min at a time, on at least 3 days a week (1 – 2 years). The proportion who met this guideline at 2 years of age was too high to allow estimation of mean differences (98.0%). Not measured at 3.5 years

<sup>g</sup> 11 – 14 h (1 – 2 years) or 10 – 13 h (3 – 5 years). The proportion who met this guideline at 2 and 3.5 years of age and at 2 or more time points was too high to allow estimation of mean differences (92.3, 95.5, and 92.5% respectively)

<sup>h</sup> Having a consistent sleep time or consistent wake time defined as within-person standard deviation of < 30 min

<sup>i</sup> The proportion who met this guideline at two or more time points was too low to allow estimation of mean differences (9.8%)

Bold figures denote  $P < 0.05$

cross-sectional study in 4-year old children reported that adherence to all three guidelines was marginally associated with better Theory of Mind performance, a measure of a child's ability to attribute core mental states, but had no effect on emotional understanding as measured using the Test of Emotion Comprehension [15]. Another cross-sectional study in 3-year old children determined that while children meeting more guidelines did have lower scores for emotional and behavioral problems as measured using the Child Behavior Checklist, the observed effect size was small [16]. Only two longitudinal studies appear to have been undertaken, both demonstrating no evidence of association between compliance to individual or combined guidelines and a variety of psychosocial outcomes one [18] to three [17] years later. These findings fit with the mixed body of literature examining each individual behavior in relation to outcomes of interest. Although previous reviews have demonstrated that children with sleep problems are more likely to develop emotional and behavioral problems such as depression and anxiety [3, 41, 42], compliance analyses such as in the current study focus on sleep duration (and in our case also sleep regularity in terms of sleep and wake timings) rather than examining the presence or otherwise of sleep problems. In our study, scores for anxiety were lower in those who met guidelines for sleep duration at one year of age (11–14 h a night), scores for resilience were higher for those with consistent wake schedules at one year of age, and children who met both types of sleep guideline at 5 years of age showed improvements in inhibitory control. However, significant differences were not observed at any other ages or measures examined, illustrating limited consistency in the relationships observed. Others have also reported no relationship between meeting sleep guidelines and inhibition [18] or measures of externalizing and internalizing problems [16, 18] or emotional skills or prosocial behaviors scores [17], although Cliff et al. [15] reported significantly greater emotional comprehension in children who met sleep duration guidelines.

Similarly, while previous reviews have demonstrated associations between physical activity and mental health in children, the effect sizes are often very small, the research designs are often weak, and considerable heterogeneity exists [4, 43]. For example, none of the these previous 24-h movement guidelines papers found any significant relationship between meeting activity guidelines and any measurement of mental or psychosocial health examined [15–18]. A recent review in older children also demonstrated that relationships between physical activity and mental health outcomes were less apparent than those observed for sleep and screen time compliance, although the quality of the evidence was very low, with all studies examined being cross-sectional in design [40].

By contrast, our findings regarding the benefits of restricting screen time on reducing levels of depression and anxiety just prior to starting school did appear more consistent, with potentially meaningful effect sizes observed at many of the time points examined. Beneficial effects were particularly evident in children who met guidelines at two or more time points. None of the previous 24-h movement papers specifically examined anxiety or depression, with most reporting no significant relationships with a variety of other measures [15, 17, 18], with the exception of Carson et al. [16] who reported higher externalizing and internalizing problem scores in children who did not meet screen time guidelines. Previous reviews have also indicated that the relationships between screen time and psychosocial outcomes are inconsistent in young children, with few measuring anxiety or depression for comparison [44]. However, a recent review of reviews has highlighted that there is moderately strong evidence for an adverse association between screen time and depressive symptoms (with a weak association for anxiety) in children and adolescents in general [45]. Why our study observed more consistent findings for screen time compared with physical activity or sleep is not clear, particularly given the parent-reported nature of our screen time data in comparison with data on sleep and physical activity which were objectively measured. Unfortunately, little is currently known regarding the underlying mechanisms explaining how and why sleep and activity may be beneficial, and screen time not, for mental health in children [4]. Few studies have examined potential mechanisms and a high level of study heterogeneity exists, limiting firm conclusions.

It is clear that many questions remain about the relationships between sleep, physical activity, and screen time in relation to mental health and wellbeing, with a strong need for prospective, high-quality studies that use robust measures of all movement behaviors and validated measures of mental health to increase our understanding in this topic area [40]. Repeated measures of key behaviors may help address issues relating to the heterogeneity of findings at this young age, given the rapid changes in development that occur both behaviourally and emotionally. Psychosocial outcomes need to be examined in a multitude of ways, including both subjective and objective tests where possible, given the relative strengths and weaknesses of each. It is also feasible that the mixed findings to date arise as a result of insufficient consideration of whether any critical windows exist for 24-h movement behaviors being able to influence later mental health and wellbeing in young children.

Our study has several strengths including the assessment of both sleep and physical activity by actigraphy, over several days, and at multiple time points, and our

evaluation of the second 'level' of sleep (consistency) and sedentary (restraint) guidelines which have rarely been examined to date. We also evaluated multiple aspects of mental health and wellbeing, using both parental report assessing 'usual' behavior, and a more 'moment-in-time' lab-based assessment by trained measurers. Measuring children's self-regulation skills via laboratory tasks represents a significant strength of this study. However, our study also has some limitations. While our measurement of screen time covered multiple types of device use, reflecting the rapid changes in technology use over recent times, we did not collect quantifiable data at 3.5 years of age limiting any comparison against guidelines at this time. Although our sample size was reasonably large, we did not have sufficient participant numbers to investigate whether differences in the relationship between adherence to guidelines and mental health and wellbeing outcomes were apparent in boys and girls. We also could not assess the relationships at several of the time points because of ceiling effects with very high numbers of children achieving the requisite guideline (physical activity at 2 years, sleep at 2 and 3.5 years). It is perhaps surprising that such high numbers of children achieved the sleep guidelines given that we measured sleep using actigraphy, which typically provides values for sleep duration that are shorter than those obtained by questionnaire [46]. However, current Canadian sleep guidelines have been predominantly developed on the basis of questionnaire data on sleep duration [11], meaning that it is likely we have underestimated the number of children who met sleep guidelines. In the absence of alternatives, we used accelerometer cut-points that were developed for older toddlers [32, 33] than the youngest age point included in our sample (3 years versus 1 year) and the same cut points at each age point. As children's motor development changes rapidly during the first two years of life, the use of these cut-points may have introduced measurement inaccuracy. Finally, multiple measures of mental health and psychosocial functioning were presented, but these were mostly obtained from a single questionnaire, the BASC-2, in which some items are used in a number of different subscales. However, while the different constructs represented by the subscales might be correlated, the subscales were examined in separate models.

## Conclusions

Although adherence to some individual movement guidelines at certain ages throughout early childhood was associated with improved mental health and wellbeing at 5 years of age, particularly reduced anxiety and depression scores, there was little consistency in these relationships. Therefore, it is not possible to conclude that better mental health and wellbeing outcomes result

for pre-schoolers meeting any one, or all movement guidelines from this research. Perhaps a different analytic approach is required such as examining compositional time use [47, 48] in relation to mental wellbeing in children. Compositional analyses, a relatively recent phenomenon, investigate all movement behaviours (sleep, activity, sedentary time) simultaneously, by accounting for the finite nature of the 24-h window, where if time spent in one component increases (e.g. sleep), then another component within the same 24-h block *must* decrease (e.g. sedentary time or physical activity). While researchers are starting to examine compositional time use in relation to mental wellbeing in children cross-sectionally [49, 50], longitudinal data are scarce [51].

## Abbreviations

LMVPA: Light-to-vigorous physical activity; MVPA: Moderate-to-vigorous physical activity; BASC-2: Behavioral Assessment System for Children; NEPSY-2: NEuroPSYchological Assessment.

## Acknowledgements

We are grateful to all the families who participated in the POI study, and to the POI research staff in the Departments of Women's and Children's Health, and Medicine at the University of Otago.

## Authors' contributions

RWT and BJT are Principal Investigators of POI and designed the overall project along with DH, KM-J and BCG; RWT conceived the idea for this study; KMJ undertook the accelerometry analyses; JJH designed and undertook the statistical analyses; RWT wrote the first and subsequent drafts of the manuscript and all authors critically revised the manuscript for important intellectual content. RWT had primary responsibility for final content. All authors read and approved the final manuscript.

## Funding

The POI Study was funded by the Health Research Council of New Zealand (08/374, 12/281, 12/310) and the Southern District Health Board. RWT is supported by a Fellowship from Karitane Products Society. The funding source had no role in the design and conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or the decision to submit the manuscript for publication.

## Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

The original intervention was approved by the Lower South Ethics Committee (LRS/12/08/063) and the follow-up study by the University of Otago Human Ethics Committee (12/274). Written informed consent was obtained from the parent/guardian of all child participants.

### Consent for publication

Not applicable.

### Competing interests

The authors declare they have no competing interests.

### Author details

<sup>1</sup>Department of Medicine, University of Otago, PO Box 56, Dunedin, New Zealand. <sup>2</sup>Biostatistics Centre, University of Otago, Dunedin, New Zealand. <sup>3</sup>Department of Psychology, University of Otago, Dunedin, New Zealand.

<sup>4</sup>Department of Women's and Children's Health, University of Otago, Dunedin, New Zealand.

Received: 19 May 2021 Accepted: 10 August 2021

Published online: 25 August 2021

## References

- Mental health: a state of well-being [[http://www.who.int/features/factfiles/mental\\_health/en/](http://www.who.int/features/factfiles/mental_health/en/)]. Accessed 5 Apr 2021.
- Robson DA, Allen MS, Howard SJ. Self-regulation in childhood as a predictor of future outcomes: a meta-analytic review. *Psychol Bull.* 2020;146:324–54.
- Chaput JP, Gray CE, Poitras VJ, Carson V, Gruber R, Olds TS, Weiss SK, Gorber SC, Kho ME, Sampson M, et al. Systematic review of the relationships between sleep duration and health indicators in school-aged children and youth. *Appl Physiol Nutr Metab.* 2016;41:5266–82.
- Lubans D, Richards J, Hillman C, Faulkner G, Beauchamp M, Nilsson M, Kelly P, Smith J, Raine L, Biddle S. Physical activity for cognitive and mental health in youth: a systematic review of mechanisms. *Pediatrics.* 2016;138:e20161642.
- Tremblay MS, LeBlanc AG, Kho ME, Saunders TJ, Larouche R, Colley RC, Goldfield G, Connor GS. Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act.* 2011;8:98.
- Chaput JP, Gray CE, Poitras VJ, Carson V, Gruber R, Birken CS, MacLean JE, Aubert S, Sampson M, Tremblay MS. Systematic review of the relationships between sleep duration and health indicators in the early years (0–4 years). *BMC Pub Health.* 2017;17:855.
- Carson V, Lee E-Y, Hewitt L, Jennings C, Hunter S, Kuzik N, Stearns JA, Unrau SP, Poitras VJ, Gray C, et al. Systematic review of the relationships between physical activity and health indicators in the early years (aged 0–4 years). *BMC Pub Health.* 2017;17:854.
- Poitras VJ, Gray CE, Janssen X, Aubert S, Carson V, Faulkner G, Goldfield GS, Reilly JJ, Sampson M, Tremblay MS. Systematic review of the relationships between sedentary behaviour and health indicators in the early years (0–4 years). *BMC Pub Health.* 2017;17:868.
- Williams SM, Taylor BJ, Taylor RW. Do more active children sleep more? A repeated cross-sectional analysis using accelerometry. *PLoS One.* 2014;9:e93117.
- Rollo S, Antsygina O, Tremblay MS. The whole day matters: understanding 24-hour movement guideline adherence and relationships with health indicators across the lifespan. *J Sport Health Sci.* 2020;9:493–510.
- Tremblay MS, Carson V, Chaput JP, Adamo KB, Aubert S, Choquette L, et al. Canadian 24-hour movement guidelines for the early years (0–4 years): an integration of physical activity, sedentary behavior, and sleep. *BMC Pub Health.* 2017;17:874.
- Australian 24-hour movement guidelines for the early years (birth to 5 years): an integration of physical activity, sedentary behaviour, and sleep [<https://www1.health.gov.au/internet/main/publishing.nsf/Content/npra-0-5yrs-brochure>]. Accessed 14 Apr 2021.
- Sit less, move more, sleep well: active play guidelines for under-fives [<https://www.health.govt.nz/publication/sit-less-move-more-sleep-well-active-play-guidelines-under-fives>]. Accessed 14 Apr 2021.
- Guidelines on physical activity, sedentary behaviour and sleep for children under 5 years of age [<https://apps.who.int/iris/handle/10665/311664>]. Accessed 14 Apr 2021.
- Cliff DP, McNeill J, Vella SA, Howard SJ, Santos R, Batterham M, Melhuish E, Okely AD, de Rosnay M. Adherence to 24-hour movement guidelines for the early years and associations with social-cognitive development among Australian preschool children. *BMC Pub Health.* 2017;17:857.
- Carson V, Ezeugwu VE, Tamana SK, Chjikuma J, Lefebvre DL, Azad MB, Moraes TJ, Subbarao P, Becker AB, Turvey SE, et al. Associations between meeting the Canadian 24-hour movement guidelines for the early years and behavioral and emotional problems among 3-year-olds. *J Sci Med Sport.* 2019;22:797–802.
- Hinkley T, Timperio A, Watson A, Duckham RL, Okely AD, Cliff DP, Carver A, Hesketh KD. Prospective associations with physiological, psychosocial and educational outcomes of meeting Australian 24-hour movement guidelines for the early years. *Int J Behav Nutr Phys Act.* 2020;17:36.
- McNeill J, Howard SJ, Vella SA, Cliff DP. Compliance with the 24-Hour movement guidelines for the early years: Cross-sectional and longitudinal associations with executive function and psychosocial health in preschool children. *J Sci Med Sport.* 2020;23:846–53.
- Chia MYH, Tay LY, Chua TBK. Quality of life and meeting 24-h WHO guidelines among preschool children in Singapore. *Early Child Educ J.* 2020;48:313–23.
- Meredith-Jones K, Haszard JJ, Moir C, Heath A-L, Galland BC, Taylor BJ, Gray AR, Sayers RM, Taylor RW. Physical activity and inactivity trajectories associated with body composition in preschoolers. *Int J Obes.* 2018;42:1621–30.
- Taylor BJ, Heath A-L, Galland BC, Gray AR, Lawrence JA, Sayers RM, Dale K, Coppell KJ, Taylor RW. Prevention of overweight in infancy (POI.nz) study: a randomised controlled trial of sleep, food and activity interventions for preventing overweight from birth. *BMC Pub Health.* 2011;11:942.
- Taylor RW, Heath A-L, Galland BC, Cameron SL, Lawrence JA, Gray AR, Tannock GW, Lawley B, Healey D, Sayers RM, et al. Three-year follow-up of a randomised controlled trial to reduce excessive weight gain in the first two years of life: protocol for the POI follow-up study. *BMC Pub Health.* 2016;16:771.
- Moir C, Meredith-Jones K, Taylor BJ, Gray AR, Heath ALM, Dale K, Galland BC, Lawrence JA, Sayers RM, Taylor RW. Early intervention to encourage physical activity in infants and toddlers: an RCT. *Med Sci Sports Exerc.* 2016;48:2246–453.
- Taylor BJ, Gray AR, Galland BC, Heath A-L, Lawrence JA, Sayers RM, Cameron SL, Hanna M, Dale K, Coppell KJ, Taylor RW. Targeting sleep, food, and activity in infants for obesity prevention: an RCT. *Pediatrics.* 2017;139:e20162037.
- Taylor RW, Gray AR, Heath A-L, Galland BC, Lawrence JA, Sayers RM, Healey D, Tannock GW, Meredith-Jones K, Hanna M, et al. Sleep, nutrition and physical activity interventions to prevent obesity in infancy: Follow-up of the POI randomized controlled trial at 3.5 & 5 years. *Am J Clin Nutr.* 2018;108:228–36.
- Atkinson J, Salmond C, Crampton P. NZDep2013 index of deprivation. Department of Public Health: University of Otago. 2014;5541:1–64.
- de Onis M, Onyango AW, Van den Broeck J, Chumlea WC, Martorell R. Measurement and standardization protocols for anthropometry used in the construction of a new international growth reference. *Food Nutr Bull.* 2004;25:S27–36.
- World Health Organization. WHO child growth standards based on length/height, weight and age. *Acta Paediatr Suppl.* 2006;450:76–85.
- Galland BC, Kennedy GJ, Mitchell RA, Taylor BJ. Algorithms for using an activity-based accelerometer for identification of infant sleep-wake states during nap studies. *Sleep Med.* 2012;13:743–51.
- Meredith-Jones K, Williams SM, Galland BC, Kennedy G, Taylor RW. 24hr accelerometry: Impact of sleep-screening methods on estimates of physical activity and sedentary time. *J Sport Sci.* 2016;34:679–85.
- Esliger DW, Copeland JL, Barnes JD, Tremblay MS. Standardizing and optimizing the use of accelerometer data for free-living physical activity monitoring. *J Phys Act Health.* 2005;2:366–83.
- Adolph A, Puyau M, Vohra F, Nicklas T, Zakeri I, Butte NF. Validation of uniaxial and triaxial accelerometers for the assessment of physical activity in children. *J Phys Act Health.* 2012;9:944–53.
- Janssen X, Cliff DP, Reilly JJ, Hinkley T, Jones R, Batterham M, Ekelund U, Brage S, Okely AD. Evaluation of actical equations and thresholds to predict physical activity intensity in young children. *Phys Act Health.* 2015;33:498–506.
- Reynolds CR, Kamphaus RW. *BASC-2: behavior assessment system for children.* 2nd ed. Bloomington: Pearson Assessments; 2004.
- Korkman M, Kirk U, Kemp S. *NEPSY - second edition (NEPSY - II).* San Antonio: Pearson Education; 2007.
- McClelland MM, Cameron CE. Self-regulation in early childhood: Improving conceptual clarity and developing ecologically valid measures. *Child Dev Perspect.* 2012;6:136–42.
- Greenland S, Senn SJ, Rothman KJ. Statistical tests, P values, confidence intervals, and power: a guide to misinterpretations. *Europ J Epidemiol.* 2016;31:337–50.
- Schafer T, Schwarz MA. The meaningfulness of effect sizes in psychological research: Differences between sub-disciplines and the impact of potential biases. *Front Psychol.* 2019. <https://doi.org/10.3389/fpsyg.2019.00813>.

39. Meredith-Jones KA, Taylor RW, Haszard JJ, Galland BC, Heath A-LM, Lawrence J, Gray AR, Sayers RC, Hanna M, Hatch B, Taylor BJ. Do young children consistently meet 24-hour movement guidelines? A longitudinal analysis using actigraphy. *Int J Obes*. 2019;43:2555–64.
40. Sampasa-Kanyinga H, Colman I, Goldfield G, Janssen I, Wang J, Podinic I, Tremblay MS, Saunders TJ, Sampson M, Chaput JP. Combinations of physical activity, sedentary time, and sleep duration and their association with depressive symptoms and other mental health problems in children and adolescents: a systematic review. *Int J Behav Nutr Phys Act*. 2020;17:72.
41. Sadeh A, Liat T, Kahn M. Sleep in infancy and childhood: implication for emotional and behavioral difficulties in adolescence and beyond. *Curr Opin Psychiatr*. 2014;27:453–9.
42. McMakin DL, Alfano CA. Sleep and anxiety in late childhood and early adolescence. *Curr Opin Psychiatr*. 2015;28:483–9.
43. Biddle SJH, Asare M. Physical activity and mental health in children and adolescents: a review of reviews. *Brit J Sports Med*. 2011;45:886–95.
44. Hinkley T, Teychenne M, Downing KL, Ball K, Salmon J, Hesketh KD. Early physical activity, sedentary behaviors and psychosocial wellbeing: a systematic review. *Prev Med*. 2014;62:182–92.
45. Stiglic N, Viner RM. Effects of screentime on the health and well-being of children and adolescents: a systematic review of reviews. *BMJ Open*. 2019;9:e023191.
46. Galland BC, Short MA, Terrill P, Rigney G, Haszard JJ, Coussens S, Foster-Owens M, Biggs S. Establishing normative values for pediatric nighttime sleep measured by actigraphy: a systematic review and meta-analysis. *Sleep*. 2018;41:1–16. <https://doi.org/10.1093/sleep/zsy1017>.
47. Pedišić Z. Measurement issues and poor adjustments for physical activity and sleep undermine sedentary behaviour research—the focus should shift to the balance between sleep, sedentary behaviour, standing and activity. *Kinesiology*. 2014;46:135–46.
48. Kuzik N, Poitras VJ, Tremblay MS, Lee E-Y, Hunter S, Carson V. Systematic review of the relationships between combinations of movement behaviours and health indicators in the early years (0–4 years). *BMC Pub Health*. 2017;17:849.
49. Bezerra TA, Clark CCT, Filho ANDS, Fortes LDS, Mota JAPS, Duncan MJ, Martins CMDL. 24-hour movement behaviour and executive function in preschoolers: A compositional and isotemporal reallocation analysis. *Europ J Sport Sci*. 2021;21:1064–72.
50. Fairclough SJ, Tyler R, Dainty JR, Dumuid D, Richardson C, Shepstone L, Atkin AJ. Cross-sectional associations between 24-hour activity behaviours and mental health indicators in children and adolescents: A compositional data analysis. *J Sports Sci*. 2021;39:1602–14.
51. Chong KH, Parrish A-M, Okely AD. Cross-sectional and longitudinal associations between 24-hour movement behaviours, recreational screen use and psychosocial health outcomes in children: A compositional data analysis approach. *Int J Environ Res Public Health*. 2021;18:0603.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

